

Appendix 3.

Hospitalisations by State/Territory and financial year (July 1997–June 2002)

Table 27. Hospitalisations by State/Territory and financial year (July 1997–June 2002)

Disease*	Year	Number of hospitalisations										Hospitalisation rate per 100,000 population									
		ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total		
Diphtheria†	97/98	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	98/99	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	99/00	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	
	00/01	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	01/02	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Total†	0	2	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	
<i>Haemophilus influenzae</i> type b disease (<15 yr only)	97/98	1	34	3	18	4	2	13	43	118	1.5	2.6	6.0	2.4	1.3	1.9	1.4	10.9	3.0		
	98/99	1	17	1	15	6	2	8	10	60	1.5	1.3	2.0	2.0	2.0	2.0	0.8	2.5	1.5		
	99/00	1	9	1	18	3	1	11	5	49	1.5	0.7	2.0	2.4	1.0	1.0	1.2	1.3	1.2		
	00/01	0	4	1	23	5	4	8	2	47	0.0	0.3	2.0	3.0	1.7	4.0	0.8	0.5	1.2		
	01/02	0	17	3	9	4	0	6	5	44	0.0	1.3	5.9	1.2	1.4	0.0	0.6	1.2	1.1		
	Total†	3	81	9	83	22	9	46	65	318	0.9	1.2	3.6	2.2	1.5	1.8	1.0	3.3	1.6		
Hepatitis A	97/98	11	393	20	291	53	0	96	74	938	3.6	6.3	10.7	8.6	3.6	0.0	2.1	4.1	5.1		
	98/99	5	256	11	252	64	4	110	74	781	1.6	4.0	5.8	7.3	4.3	0.8	2.4	4.1	4.2		
	99/00	1	181	20	122	57	6	157	100	649	0.3	2.8	10.4	3.5	3.8	1.3	3.4	5.4	3.4		
	00/01	5	120	13	66	28	7	75	32	346	1.6	1.9	6.6	1.9	1.9	1.5	1.6	1.7	1.8		
	01/02	5	145	14	60	28	3	54	16	325	1.6	2.2	7.1	1.7	1.9	0.6	1.1	0.8	1.7		
	Total†	27	1,095	78	791	230	20	492	296	3,039	1.7	3.4	8.1	4.5	3.1	0.8	2.1	3.2	3.2		
Hepatitis B (acute) (principal diagnosis only)	97/98	3	54	1	23	13	0	66	15	175	1.0	0.9	0.5	0.7	0.9	–	1.4	0.8	0.9		
	98/99	2	45	3	31	12	0	70	20	187	0.6	0.7	1.6	0.9	0.8	–	1.5	1.1	1.0		
	99/00	1	46	3	19	11	1	53	21	158	0.3	0.7	1.6	0.5	0.7	0.2	1.1	1.1	0.8		
	00/01	0	36	1	14	10	10	65	22	158	–	0.6	0.5	0.4	0.7	2.1	1.4	1.2	0.8		
	01/02	0	46	6	13	4	1	61	16	147	–	0.7	3.0	0.4	0.3	0.2	1.3	0.8	0.8		
	Total†	6	227	14	100	50	12	315	94	825	0.4	0.7	1.5	0.6	0.7	0.5	1.3	1.0	0.9		

* Refer to Chapter 3 for case definitions.

† Total cases for 5-year period and average annual rate per 100,000 population.

‡ Hospitalisations for rare diseases such as diphtheria, poliomyelitis and tetanus should be interpreted with extreme caution due to possible coding errors. For NSW, polio, tetanus and diphtheria occurrences are considered inaccurate, given no notifications of these diseases have been identified to NSW Health for the period 2001–2002. Caution should be exercised in analysing these data and users should consult the disease notifications as published in the *New South Wales Public Health Bulletin*.

Table 27. Hospitalisations by State/Territory and financial year (July 1997–June 2002)

Disease*	Year	Number of hospitalisations							Total	Hospitalisation rate per 100,000 population									
		ACT	NSW	NT	Qld	SA	Tas	Vic		WA	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total
Influenza	97/98	37	1,698	32	1,058	559	68	1,591	1,090	6,133	12.0	27.1	17.1	31.2	37.7	14.4	34.6	60.7	33.1
	98/99	20	1,494	46	1,145	442	36	836	573	4,592	6.5	23.6	24.2	33.2	29.7	7.6	18.0	31.4	24.5
	99/00	10	1,154	25	858	282	41	781	847	3,998	3.2	18.0	13.0	24.5	18.8	8.7	16.7	45.8	21.1
	00/01	17	1,190	38	700	331	54	602	536	3,468	5.4	18.3	19.4	19.7	22.0	11.5	12.7	28.6	18.1
	01/02	11	817	24	631	293	36	493	502	2,807	3.4	12.4	12.1	17.4	19.4	7.6	10.3	26.4	14.5
	Total†	95	6,353	165	4,392	1,907	235	4,303	3,548	20,998	6.1	19.8	17.1	25.0	25.5	10.0	18.3	38.4	22.2
Measles	97/98	11	63	1	59	4	3	12	3	156	3.6	1.0	0.5	1.7	0.3	0.6	0.3	0.2	0.8
	98/99	0	17	0	13	6	3	30	8	78	–	0.3	–	0.4	0.4	0.6	0.6	0.4	0.4
	99/00	1	9	14	10	3	1	15	7	67	0.3	0.1	7.3	0.3	0.2	0.2	0.3	0.4	0.4
	00/01	2	13	0	6	4	0	36	3	64	0.6	0.2	–	0.2	0.3	–	0.8	0.2	0.3
	01/02	0	10	0	2	1	2	18	8	41	–	0.2	–	0.1	0.1	0.4	0.4	0.4	0.2
	Total†	14	112	15	90	18	9	111	29	406	0.9	0.3	1.6	0.5	0.2	0.4	0.5	0.3	0.4
Meningococcal disease	97/98	6	335	17	179	33	14	102	75	761	1.9	5.3	9.1	5.3	2.2	3.0	2.2	4.2	4.1
	98/99	2	284	15	155	39	23	105	124	749	0.6	4.5	7.9	4.5	2.6	4.9	2.3	6.8	4.0
	99/00	5	297	13	108	51	18	210	107	817	1.6	4.6	6.7	3.1	3.4	3.8	4.5	5.8	4.3
	00/01	5	320	10	140	51	15	230	96	867	1.6	4.9	5.1	3.9	3.4	3.2	4.9	5.1	4.5
	01/02	9	240	15	178	60	44	253	77	876	2.8	3.7	7.6	4.9	4.0	9.3	5.3	4.1	4.5
	Total†	27	1,476	70	760	324	114	900	479	4,070	1.7	4.6	7.3	4.3	3.1	4.8	3.8	5.2	4.3
Mumps	97/98	2	18	1	10	3	1	8	8	51	0.6	0.3	0.5	0.3	0.2	0.2	0.2	0.4	0.3
	98/99	1	35	0	9	6	1	4	2	85	0.3	0.6	–	0.3	0.4	0.2	0.1	0.1	0.3
	99/00	2	22	0	14	2	0	7	7	54	0.6	0.3	–	0.4	0.1	–	0.1	0.4	0.3
	00/01	3	15	0	10	4	0	12	4	48	1.0	0.2	–	0.3	0.3	–	0.3	0.2	0.3
	01/02	0	18	0	5	4	1	7	2	37	–	0.3	–	0.1	0.3	0.2	0.1	0.1	0.2
	Total†	8	108	1	48	19	3	38	23	248	0.5	0.3	0.1	0.3	0.3	0.1	0.2	0.2	0.3
Pertussis	97/98	9	439	8	213	123	11	125	237	1,165	2.9	7.0	4.3	6.3	8.3	2.3	2.7	13.2	6.3
	98/99	2	171	2	87	25	1	79	27	396	0.6	2.7	1.1	2.5	1.7	0.2	1.7	1.5	2.1
	99/00	8	115	2	53	13	21	118	17	349	2.6	1.8	1.0	1.5	0.9	4.5	2.5	0.9	1.8
	00/01	8	269	10	55	78	6	62	19	507	2.5	4.1	5.1	1.5	5.2	1.3	1.3	1.0	2.6
	01/02	4	260	44	156	134	6	98	68	770	1.3	4.0	22.2	4.3	8.9	1.3	2.0	3.6	4.0
	Total†	31	1,254	66	564	373	45	482	368	3,187	2.0	3.9	6.9	3.2	5.0	1.9	2.1	4.0	3.4
(<5yr only)	Total†	22	884	57	362	213	25	341	295	2,200	20.7	40.4	64.6	29.6	45.1	15.6	21.9	46.3	34.2

* Refer to Chapter 3 for case definitions.

† Total cases for 5-year period and average annual rate per 100,000 population.

Table 27. Hospitalisations by State/Territory and financial year (July 1997–June 2002) (continued)

Disease*	Year	Number of hospitalisations							Hospitalisation rate per 100,000 population										
		ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total
Pneumococcal disease (invasive) [†]	97/98	6	260	32	71	63	12	143	54	641	1.9	4.1	17.1	2.1	4.3	2.5	3.1	3.0	3.5
	98/99	9	292	54	88	74	16	173	44	753	2.9	4.6	28.4	2.6	5.0	3.4	3.7	2.4	4.0
	99/00	27	333	43	116	82	23	200	118	949	8.6	5.2	22.3	3.3	5.5	4.9	4.3	6.4	5.0
	00/01	14	373	58	170	65	13	218	99	1,010	4.4	5.8	29.7	4.8	4.3	2.8	4.6	5.3	5.3
	01/02	20	365	46	189	104	25	254	98	1,101	6.3	5.6	23.3	5.2	6.9	5.3	5.3	5.2	5.7
	Total [†]	76	1,623	233	634	388	89	988	413	4,454	4.9	5.1	24.2	3.6	5.2	3.8	4.2	4.5	4.7
Poliomyelitis [§]	97/98	0	31	0	19	6	1	14	0	71	–	0.5	–	0.6	0.4	0.2	0.3	–	0.4
	98/99	0	34	0	11	1	2	20	1	69	–	0.5	–	0.3	0.1	0.4	0.4	0.1	0.4
	99/00	0	1	0	0	4	0	16	0	21	–	0.0	–	–	0.3	–	0.3	–	0.1
	00/01	0	15	1	0	3	0	1	0	20	–	0.2	0.5	–	0.2	–	0.0	–	0.1
	01/02	1	6	3	0	2	2	0	0	14	0.3	0.1	1.5	–	0.1	0.4	–	–	0.1
	Total [†]	1	87	4	30	16	5	51	1	195	0.1	0.3	0.4	0.2	0.2	0.2	0.2	0.0	0.2
(Principal diagnosis only)	Total [†]	0	3	0	2	1	0	2	0	8	–	0.0	–	0.0	0.0	–	0.0	–	0.0
	97/98	1	22	0	12	5	0	3	5	48	0.3	0.4	–	0.4	0.3	–	0.1	0.3	0.3
	98/99	0	16	0	9	3	1	11	3	43	–	0.3	–	0.3	0.2	0.2	0.2	0.2	0.2
	99/00	0	8	0	8	3	0	10	0	29	–	0.1	–	0.2	0.2	–	0.2	–	0.2
	00/01	0	24	0	3	1	0	4	1	33	–	0.4	–	0.1	0.1	–	0.1	0.1	0.2
	01/02	0	15	0	1	1	1	3	0	21	–	0.2	–	0.0	0.1	0.2	0.1	–	0.1
Total [†]	1	85	0	33	13	2	31	9	174	0.1	0.3	–	0.2	0.2	0.1	0.1	0.1	0.2	

* Refer to Chapter 3 for case definitions.

† Total cases for 5-year period and average annual rate per 100,000 population.

‡ Pneumococcal meningitis and septicaemia.

§ Hospitalisations for rare diseases such as diphtheria, poliomyelitis and tetanus should be interpreted with extreme caution due to possible coding errors. For NSW, polio, tetanus and diphtheria occurrences are considered inaccurate, given no notifications of these diseases have been identified to NSW Health for the period 2001–2002. Caution should be exercised in analysing these data and users should consult the disease notifications as published in the *New South Wales Public Health Bulletin*.

Table 27. Hospitalisations by State/Territory and financial year (July 1997–June 2002) (continued)

Disease*	Year	Number of hospitalisations								Hospitalisation rate per 100,000 population										
		ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total	
Tetanus†	97/98	0	20	0	2	3	1	4	4	34	–	0.3	–	0.1	0.2	0.2	0.1	0.1	0.2	0.2
	98/99	0	15	0	6	0	1	4	7	33	–	0.2	–	0.2	–	0.2	0.1	0.4	0.2	0.2
	99/00	0	8	0	8	6	2	5	2	32	–	0.1	–	0.2	0.4	0.4	0.1	0.1	0.2	0.2
	00/01	0	17	0	6	3	1	3	1	31	–	0.3	–	0.2	0.2	0.2	0.1	0.1	0.1	0.2
	01/02	0	6	0	4	3	0	4	6	23	–	0.1	–	0.1	0.2	0.0	0.1	0.3	0.1	0.1
	Total‡	0	66	0	26	15	5	20	20	153	–	0.2	–	0.1	0.2	0.2	0.1	0.2	0.1	0.2
Varicella	97/98	27	440	18	300	126	25	365	214	1,515	8.7	7.0	9.6	8.8	8.5	5.3	7.9	11.9	8.2	
	98/99	22	677	33	427	188	36	399	199	1,991	7.1	10.7	17.4	12.4	12.6	7.6	8.6	10.9	10.6	
	99/00	30	580	32	332	93	48	418	190	1,734	9.6	9.0	16.6	9.5	6.2	10.2	8.9	10.3	9.2	
	00/01	22	544	14	335	152	30	344	198	1,639	7.0	8.4	7.2	9.4	10.1	6.4	7.3	10.6	8.6	
	01/02	22	570	21	308	185	32	344	197	1,679	6.9	8.7	10.6	8.5	12.2	6.8	7.2	10.4	8.6	
	Total‡	123	2,811	118	1,702	744	171	1,870	998	8,558	7.9	8.8	12.3	9.7	9.9	7.2	8.0	10.8	9.0	
Zoster	97/98	37	1,496	22	776	486	115	1,042	421	4,395	12.0	23.8	11.8	22.9	32.8	24.3	22.7	23.5	23.7	
	98/99	33	1,717	21	808	508	123	1,074	422	4,718	10.6	27.1	11.1	23.4	34.1	26.1	23.2	23.2	25.2	
	99/00	35	1,574	26	853	519	147	1,103	406	4,673	11.2	24.6	13.5	24.4	34.7	31.2	23.5	21.9	24.7	
	00/01	44	1,499	15	838	444	135	1,142	416	4,533	14.0	23.1	7.7	23.5	29.5	28.6	24.1	22.2	23.7	
	01/02	56	1,453	21	891	514	138	1,175	380	4,628	17.5	22.1	10.6	24.6	34.0	29.2	24.5	20.0	23.8	
	Total‡	205	7,739	105	4,166	2,471	658	5,536	2,045	22,947	13.1	24.1	10.9	23.8	33.0	27.9	23.6	22.1	24.2	

* Refer to Chapter 3 for case definitions.

† Total cases for 5-year period and average annual rate per 100,000 population.

‡ Hospitalisations for rare diseases such as diphtheria, poliomyelitis and tetanus should be interpreted with extreme caution due to possible coding errors. For NSW, polio, tetanus and diphtheria occurrences are considered inaccurate given no notifications of these diseases have been identified to NSW Health for the period 2001–2002. Caution should be exercised in analysing these data and users should consult the disease notifications as published in the *New South Wales Public Health Bulletin*.